

RIEMANN SURFACES - SPRING 2024

EXERCISES SHEET 4

Ex 1. • Consider $\varphi: \mathbb{C} \rightarrow \mathbb{C} \xrightarrow{\pi'} \mathbb{C}/\Lambda'$. The map factors

$$z \mapsto \gamma z$$

$$w \mapsto [w]$$

through $\mathbb{C}/\Lambda$ iff $\Lambda \subseteq \text{Ker}(\varphi)$ (fundamental thm on homom.)

This is indeed the case, under the hypothesis $\gamma\Lambda \subseteq \Lambda'$:

$$z \in \Lambda \Rightarrow \varphi(z) = [\gamma z] = 0 \text{ as } \gamma z \in \Lambda'.$$

The induced map $\mathbb{C}/\Lambda \rightarrow \mathbb{C}/\Lambda'$ is simply a rescaling in bc coordinates, hence holomorphic.

Notice that φ is surjective. It is injective iff $\Lambda = \text{Ker}(\varphi)$.

We already know that $\Lambda \subseteq \text{Ker}(\varphi)$. So

$$\varphi \text{ is bijective iff } \Lambda = \text{Ker}(\varphi)$$

$$\text{iff } \gamma\Lambda = \Lambda'$$

The (unique) inverse is automatically holomorphic, induced by the map $\mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C}/\Lambda$.

$$z \mapsto \frac{z}{\gamma}$$

$$w \mapsto [w]$$

- Let $\Lambda = \{n_1\omega_1 + n_2\omega_2 \mid n_1, n_2 \in \mathbb{Z}\}$. We have two possibilities:

$$\textcircled{1} \quad \Im\left(\frac{\omega_2}{\omega_1}\right) > 0$$

$$\textcircled{2} \quad \Im\left(\frac{\omega_1}{\omega_2}\right) = \Im\left(\left(\frac{\omega_2}{\omega_1}\right)^{-1}\right) < 0$$

It cannot be zero, since ω_1 and ω_2 are linearly indep. over $\mathbb{R}$. Set $\Lambda' = \varphi\Lambda$, where

$$\begin{cases} \varphi = \frac{1}{\omega_1} & \text{in } \textcircled{1} \\ \varphi = \frac{1}{\omega_2} & \text{in } \textcircled{2} \end{cases}$$

In any case, $\Lambda' = \mathbb{Z} + \tau\mathbb{Z}$ w/ $\Im(\tau) > 0$. Besides, from the previous point, $\mathbb{C}/\Lambda \cong \mathbb{C}/\mathbb{Z} + \tau\mathbb{Z}$.

- We want to show that $\exists \varphi \in \mathbb{C}^*$ s.t. $\varphi(\mathbb{Z} + \tau'\mathbb{Z}) = \mathbb{Z} + \tau\mathbb{Z}$.

Suppose for a second that we have such φ . Then

$$\varphi \in \mathbb{Z} + \tau\mathbb{Z} \Rightarrow \exists c, d \in \mathbb{Z} \text{ s.t. } \varphi = c\tau + d$$

$$\varphi\tau' \in \mathbb{Z} + \tau\mathbb{Z} \Rightarrow \exists a, b \in \mathbb{Z} \text{ s.t. } \varphi\tau' = a\tau + b$$

so that $\tau' = \frac{a\tau + b}{c\tau + d}$. Notice that we must have

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathbb{Z})$$

for maintain linear indep, so the determinant must be ± 1 (the only invertible elem in $\mathbb{Z}$). Notice that

$$\tau' = \frac{(a\tau+b)(c\bar{\tau}+d)}{|c\tau+d|^2} = \frac{a|\tau|^2 + bd + ad\tau + bc\bar{\tau}}{|c\tau+d|^2}$$

$$\Rightarrow \Im(\tau') = \frac{ad-bc}{|c\tau+d|^2} \Im(\tau)$$

Hence, for $\tau' \in \mathbb{H}$, we must have $\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = +1$.

this cannot be zero, since
1 and τ are lin. indep./ $\mathbb{R}$.

So, we are tempted to define $\varphi = c\tau + d$. The above computations (reversed) show that indeed

$$\alpha(\mathbb{Z} + \tau'\mathbb{Z}) = \mathbb{Z} + \tau\mathbb{Z},$$

thus $T(\tau) \cong T(\tau')$.

Ex 2. We can do this in $\mathbb{C}[z_0, \dots, z_n]$.

(1 $\Rightarrow$ 2). Suppose $F \in \mathbb{C}[z_0, \dots, z_n]$. Write F as a sum of monomials:

$$F = \sum_{k \geq 0} \sum_{k_0 + \dots + k_n = k} a_{k_0, \dots, k_n} z_0^{k_0} \dots z_n^{k_n}$$

$\uparrow$ finite

Then $\forall \lambda \in \mathbb{C}^*$

$$F(\lambda z_0, \dots, \lambda z_n) = \sum_{k \geq 0} \sum_{k_0 + \dots + k_n = k} a_{k_0, \dots, k_n} (\lambda z_0)^{k_0} \dots (\lambda z_n)^{k_n}$$

$$= \sum_{k \geq 0} \lambda^k \sum_{k_0 + \dots + k_n = k} a_{k_0, \dots, k_n} z_0^{k_0} \dots z_n^{k_n}$$

This equals $\lambda^d F(z_0, \dots, z_n)$ in $\mathbb{C}[\lambda, z_0, \dots, z_n]$ iff

- for $k \neq d$, $\sum_{k_0 + \dots + k_n = k} a_{k_0, \dots, k_n} z_0^{k_0} \dots z_n^{k_n} = 0$

$$\Leftrightarrow a_{k_0, \dots, k_n} = 0 \quad \forall k_0, \dots, k_n \text{ st. } k_0 + \dots + k_n = k.$$

- for $k = d$, $F(z_0, \dots, z_n) = \sum_{k_0 + \dots + k_n = d} a_{k_0, \dots, k_n} z_0^{k_0} \dots z_n^{k_n}$

(2 $\Leftrightarrow$ 3). Suppose $F \in \mathbb{C}[z_0, \dots, z_n]$. Write F as a sum of monomials:

$$F = \sum_{k \geq 0} \sum_{k_0 + \dots + k_n = k} a_{k_0, \dots, k_n} z_0^{k_0} \dots z_n^{k_n}$$

Define the Euler operator as

$$D = \sum_{i=0}^n z_i \frac{\partial}{\partial z_i}$$

Then

$$\begin{aligned} DF(z_0, \dots, z_n) &= \sum_{k \geq 0} \sum_{k_0 + \dots + k_n = k} a_{k_0, \dots, k_n} D(z_0^{k_0} \dots z_n^{k_n}) \\ &= \sum_{k \geq 0} \sum_{k_0 + \dots + k_n = k} \underbrace{(k_0 + \dots + k_n)}_{=k} a_{k_0, \dots, k_n} z_0^{k_0} \dots z_n^{k_n} \end{aligned}$$

$$= \sum_{k \geq 0} k \sum_{k_0 + \dots + k_n = k} a_{k_0 \dots k_n} z_0^{k_0} \dots z_n^{k_n}$$

This equals d.F iff

- for $k \neq d$, $\sum_{k_0 + \dots + k_n = k} a_{k_0 \dots k_n} z_0^{k_0} \dots z_n^{k_n} = 0$

$$\Leftrightarrow a_{k_0 \dots k_n} = 0 \quad \forall k_0, \dots, k_n \text{ st. } k_0 + \dots + k_n = k.$$

- for $k = d$, $F(z_0, \dots, z_n) = \sum_{k_0 + \dots + k_n = d} a_{k_0 \dots k_n} z_0^{k_0} \dots z_n^{k_n}$

Ex 3. The eqn $x^3 + y^3 = 9$ is the dehomogenisation.

of an elliptic curve. In

particular, we have the group

law with the same defn as in

the notes, and identity


$$O = [1 : -1 : 0]$$



the homogen. curve is

$$F = z_0^3 + z_1^3 - 9z_2^3 = 0$$

and $O \in Z(F)$.



The idea is to use the

group law to produce more

rational solutions.


First, we start computing $-[(1,2) + (1,2)]$. By definition, this is the intersection of the cubic w/ the tangent at $(1,2)$.

We can do this for a generic point $P=(a,b)$ as follows.

The tangent at $P=(a,b)$ is $2(a^2x + b^2y - 9)$. Notice that $a^3 + b^3 = 9$, so the line can be written as

$$y = -\frac{a^2}{b^2}(x-a) + b.$$

The intersection w/ cubic is then



$$\begin{cases} y = -\frac{a^2}{b^2}(x-a) + b \\ x^3 + y^3 = 9 \end{cases} \rightarrow x^3 + \left(-\frac{a^2}{b^2}(x-a) + b\right)^3 = 9$$

This is a cubic eqn with a double root at $x=a$. From the theory of cubic eqns, we find that the third root is at

$$x = \frac{a(a^3 + 2b^3)}{(a^3 - b^3)} \Rightarrow y = -\frac{b(2a^3 + b^3)}{a^3 - b^3}$$

$$\text{Hence: } -2(a,b) = \left(\frac{a(a^3 + 2b^3)}{(a^3 - b^3)}, -\frac{b(2a^3 + b^3)}{a^3 - b^3} \right).$$

Notice that $(a,b) \in \mathbb{Q}^2 \Rightarrow -2(a,b) \in \mathbb{Q}^2$. Applying it to $(1,2)$, we find

$$\left(-\frac{17}{7}, \frac{20}{7} \right).$$

However, this is NOT a positive solution. The idea now is to consider the sum w/ the other solution

$$- \left[(2, 1) + \left(-\frac{17}{7}, \frac{20}{7} \right) \right].$$

which is obtained as the intersection of the line passing through them and the cubic. The line passing through the points is

$$2(13x + 31y - 57)$$

The new solution to the system

$$\begin{cases} 13x + 31y = 57 \\ x^3 + y^3 = 9 \end{cases} \quad \text{is} \quad Q = \left(-\frac{271}{438}, \frac{919}{438} \right)$$

Taking $-2Q$ gives

$$-2Q = \left(\frac{415 \ 280 \ 564 \ 497}{348 \ 671 \ 682 \ 660}, \frac{676 \ 702 \ 467 \ 503}{348 \ 671 \ 682 \ 660} \right)$$

which is now a positive rational solution!

Geometrically, the computation is pictured below.

