

RIEMANN SURFACES - SPRING 2024

EXERCISES SHEET 6

Ex 1. Let φ be a meromorphic function on $\mathbb{P}^1$. If φ is constant ($\neq \infty$), then $\varphi([z:w]) = [\text{const}: 1]$ and we are good.

If φ is non-constant, we necessarily have at least a zero and a pole (since φ is surjective), and in $\mathbb{P}^1$ -finite numbers. Let

- $x_1, \dots, x_M$ be the zeros, of order $m_1, \dots, m_M$; set $x_i = [a_i : b_i]$
- $y_1, \dots, y_N$ be the poles, of order $n_1, \dots, n_N$; set $y_j = [c_j : d_j]$.

By design, all x_i and y_j are distinct, and

$$m_1 + \dots + m_M = n_1 + \dots + n_N = \deg(f) =: d.$$

Consider now the meromorphic fct

$$\psi([z:w]) = \left[\prod_{i=1}^M (z b_i - w a_i)^{m_i} : \prod_{j=1}^N (z d_j - w c_j)^{n_j} \right]$$

It has zeros of order m_i at x_i , and poles of order n_j at y_j .

Hence, the meromorphic function φ/ψ has no zeros nor poles

(since $\text{ord}_x(\cdot)$ is a discrete valuation). We deduce that φ/ψ


is a constant ($\neq 0$ and ∞), say $[p:q] \in \mathbb{P}^1$. Hence,

$$\varphi([z:w]) = \left[\underbrace{p \cdot \prod_{i=1}^M (z b_i - w a_i)^{m_i}}_{= F(z,w)} : \underbrace{q \cdot \prod_{j=1}^N (z d_j - w c_j)^{n_j}}_{= G(z,w)} \right]$$

Both F and G are homogeneous poly, of degree d , with no common factors.

Ex 2. By the residue thm,

$$\frac{1}{2\pi i} \oint_{\partial P} f(z) dz = \sum_{\substack{z_0 \text{ pole} \\ \text{in } P}} \text{Res}_{z=z_0} f(z)$$



On the other hand,

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\partial P} f(z) dz &= \frac{1}{2\pi i} \left(\int_0^{\omega_1} - \int_{\omega_2}^{\omega_1+\omega_2} \right) f(z) dz \\ &\quad + \frac{1}{2\pi i} \left(\int_{\omega_1}^{\omega_1+\omega_2} - \int_0^{\omega_2} \right) f(z) dz \end{aligned}$$

By Λ -periodicity, the two summands are zero. Hence:

$$\sum_{\substack{z_0 \text{ pole} \\ \text{in } P}} \text{Res}_{z=z_0} f(z) = 0.$$

Now assume f has a single pole at $z_0 \in P$ that is simple. We find

$$f(z) = a_{-1} \frac{1}{z - z_0} + O(1) \quad w/ \quad a_{-1} \neq 0.$$

Since there are no other poles we find $a_{-1} = 0$, which is a contradiction.

Ex 3

i) Let K be a compact in $\mathbb{C} \setminus \Lambda$. If $z \in K$, then $|z| < R$ for some R big enough. The idea is then to split the series in two:

$$p(z) = \frac{1}{z^2} + \sum_{\substack{\omega \in \Lambda^* \\ |\omega| \leq 2R}} \left(\frac{1}{(z-\omega)^2} - \frac{1}{\omega^2} \right) + \sum_{\substack{\omega \in \Lambda^* \\ |\omega| > 2R}} \left(\frac{1}{(z-\omega)^2} - \frac{1}{\omega^2} \right)$$

The first sum is finite, and defines a continuous function on K . Hence, it is uniformly bounded:

$$\left| \frac{1}{z^2} + \sum_{\substack{\omega \in \Lambda^* \\ |\omega| \leq 2R}} \left(\frac{1}{(z-\omega)^2} - \frac{1}{\omega^2} \right) \right| \leq A \quad \forall z \in K$$

for some $A > 0$. The second sum is infinite, and it can be bounded as

$$\left| \sum_{\substack{\omega \in \Lambda^* \\ |\omega| > 2R}} \left(\frac{1}{(z-\omega)^2} - \frac{1}{\omega^2} \right) \right| \leq \sum_{\substack{\omega \in \Lambda^* \\ |\omega| > 2R}} \underbrace{\left| \frac{z(2\omega - z)}{\omega^2(z-\omega)^2} \right|}_{\leq \frac{R(2|\omega| + R)}{|\omega|^2 |\omega/2|^2} < \frac{10R}{|\omega|^3}}$$

$$\leq B \sum_{\omega \in \Lambda^*} \frac{1}{|\omega|^3}$$

where $B = 10R$. All together, we find that $\forall z \in K$

$$|\gamma(z)| \leq A + B \sum_{\omega \in \Lambda^*} \frac{1}{|\omega|^3}$$

for some $A, B > 0$. The Weierstrass M-test implies absolute and uniform convergence, provided that $\sum_{\omega \in \Lambda^*} \frac{1}{|\omega|^3} < +\infty$. To check that the sum is convergent, notice that the number of lattice points in the annulus $N \leq |z| < N+1$ grows linearly in N :

$$\#\{\omega \in \Lambda \mid N \leq |\omega| < N+1\} \leq cN \quad \text{for some } c > 0.$$

Indeed, choose $r > 0$ s.t. the balls of radius r centered at points in Λ are disjoint. Notice that

$$\begin{aligned} \#\{\omega \in \Lambda \mid N \leq |\omega| < N+1\} &\leq \frac{\text{Area}(|z| < r)}{\text{Area}(N-r \leq |z| < N+1+r)}, \end{aligned}$$

that is

$$\#\{\omega \in \Lambda \mid N \leq |\omega| < N+1\} \leq \frac{\pi((N+1+r)^2 - (N-r)^2)}{\pi r^2} \leq c \cdot N.$$

As a consequence,

$$\begin{aligned} \sum_{\omega \in \Lambda^*} \frac{1}{|\omega|^3} &\leq \sum_{N=1}^{\infty} \frac{\#\{\omega \in \Lambda \mid N \leq |\omega| < N+1\}}{N^3} \\ &\leq c \sum_{N=1}^{\infty} \frac{1}{N^2} = \frac{c\pi^2}{6} < \infty. \end{aligned}$$

ii) By absolute convergence,

$$p'(z) = -2 \sum_{w \in \Lambda} \frac{1}{(z-w)^3}$$

which is clearly Λ -periodic. It is also odd, since

$$\begin{aligned} p'(-z) &= -2 \sum_{w \in \Lambda} \frac{1}{(-z-w)^3} \\ &= +2 \sum_{w \in \Lambda} \frac{1}{(z+w)^3} \\ &= +2 \sum_{w \in \Lambda} \frac{1}{(z-w)^3} = -p'(z). \end{aligned}$$

Similarly, p is even.

To deduce Λ -periodicity of p , consider

$$f_w(z) := p(z+w) - p(z), \quad w \in \Lambda.$$

Then $f'_w(z) = p'(z+w) - p'(z) = 0$ since p' is Λ -periodic. Hence,

$$f_w(z) = \text{constant}.$$

We can determine the constant setting $z = -\frac{w}{2}$:

$$\text{constant} = f_{\omega}\left(-\frac{\omega}{2}\right) = p\left(\frac{\omega}{2}\right) - p\left(-\frac{\omega}{2}\right) = 0$$

since p is even.

We conclude that $f_{\omega} \equiv 0$, i.e. p is Λ -periodic.

iii) Notice that p' has a triple pole of order 3.

Hence, p' has 3 zeros, counted with multiplicity.

Since p' is odd, it is easy to check that

$[\frac{\omega_1}{2}]$, $[\frac{\omega_2}{2}]$, $[\frac{\omega_1 + \omega_2}{2}]$ are zeros:

$$\begin{aligned} p'\left(\frac{n_1\omega_1 + n_2\omega_2}{2}\right) &= p'\left(\frac{n_1\omega_1 + n_2\omega_2}{2} - n_1\omega_1 - n_2\omega_2\right) \\ &= p'\left(-\frac{n_1\omega_1 + n_2\omega_2}{2}\right) \\ &= -p'\left(\frac{n_1\omega_1 + n_2\omega_2}{2}\right) \end{aligned}$$

Hence, $[\frac{\omega_1}{2}]$, $[\frac{\omega_2}{2}]$, $[\frac{\omega_1 + \omega_2}{2}]$ are zeros. Since they are distinct in $\mathbb{C}/\Lambda$, there cannot be more zeros. Moreover, they are necessarily simple.

iv) A Λ -periodic meromorphic function w with double poles on Λ has Laurent expansion

$$f(z) = \frac{a}{z^2} + O(z^{-1}) \quad \text{for some } a \in \mathbb{C}.$$

Hence, $f(z) - a p(z) = O(z^{-1})$, that is, it is Λ -periodic w/ at most a simple pole on Λ . By the the Weierstrass gap thm, it is constant: $f(z) - a p(z) = b \in \mathbb{C}$.

v) We have $p(z) - \frac{1}{z^2} = \sum_{w \in \Lambda^*} \left(\frac{1}{(z-w)^2} - \frac{1}{w^2} \right)$, which evaluates zero at $z=0$. Hence, $c=0$:

$$p(z) = \frac{1}{z^2} + \frac{1}{20} g_2 z^2 + \frac{1}{28} g_3 z^4 + O(z^6)$$

Thus:

$$p'(z) = -\frac{2}{z^3} + \frac{1}{10} g_2 z + \frac{1}{7} g_3 z^3 + O(z^5)$$

We deduce that

$$(p'(z))^2 = \frac{4}{z^6} - \frac{2}{5} g_2 \frac{1}{z^2} - \frac{4}{7} g_3 + O(z)$$

$$(p(z))^3 = \frac{1}{z^6} + \frac{3}{20} g_2 \frac{1}{z^2} + \frac{3}{28} g_3 + O(z)$$

Summing up:

$$(p'(z))^2 - [4(p(z))^3 - g_2 p(z) - g_3] = O(z)$$

We deduce that the above combination is holomorphic on $\mathbb{C}/\Lambda$ and vanishes at $[0]$, hence it is constant zero.

Ex 4. We know that E_Λ is smooth, provided that

$$S := \left\{ p \in \mathbb{C}^3 \setminus \{0\} \mid F(p) = \frac{\partial F}{\partial z_0}(p) = \frac{\partial F}{\partial z_1}(p) = \frac{\partial F}{\partial z_2}(p) = 0 \right\} = \emptyset$$

$$\text{where } F = z_1^2 z_2 - z_0^3 + \frac{g_2}{4} z_0 z_2^2 + \frac{g_3}{4} z_2^3.$$

From the theory of (depressed) cubic eqns, we have that

$$x^3 - \frac{g_2}{4}x - \frac{g_3}{4}$$

has three distinct roots, provided that $-4\left(-\frac{g_2}{4}\right)^3 - 27\left(-\frac{g_3}{4}\right)^2 \neq 0$.

This is equivalent to $\Delta \neq 0$. Hence, we can write

$$F = z_1^2 z_2 - (z_0 - r_1 z_2)(z_0 - r_2 z_2)(z_0 - r_3 z_2) \quad \text{w/ } r_i \text{ distinct.}$$

Suppose now by contradiction that $\exists p = [a : b : c] \in S$.

Notice that $\frac{\partial F}{\partial z_1}(p) = 2bc = 0 \Rightarrow b = 0 \text{ or } c = 0$

If $c=0$, then $F(p) = -a^3 = 0$. Hence, $p = [0:1:0]$. But then

$$\begin{aligned} \frac{\partial F}{\partial z_2}(p) &= z_1^2 + \alpha_1(z_0 - \alpha_2 z_2)(z_0 - \alpha_3 z_2) \\ &\quad + \alpha_2(z_0 - \alpha_1 z_2)(z_0 - \alpha_3 z_2) \\ &\quad + \alpha_3(z_0 - \alpha_1 z_2)(z_0 - \alpha_2 z_2) \Big|_{[0:1:0]} \\ &= 1 \neq 0, \text{ contradiction.} \end{aligned}$$

If $b=0$, then $F(p) = (a - \alpha_1 c)(a - \alpha_2 c)(a - \alpha_3 c) = 0$, which gives $p = [\alpha_i : 0 : 1]$ for $i=1,2,3$. But then

$$\begin{aligned} \frac{\partial F}{\partial z_0}(p) &= - (z_0 - \alpha_2 z_2)(z_0 - \alpha_3 z_2) \\ &\quad - (z_0 - \alpha_1 z_2)(z_0 - \alpha_3 z_2) \\ &\quad - (z_0 - \alpha_1 z_2)(z_0 - \alpha_2 z_2) \Big|_{p=[\alpha_i:0:1]} \\ &= -(\alpha_i - \alpha_j)(\alpha_i - \alpha_k) \neq 0, \text{ contradiction.} \end{aligned}$$

$$\uparrow \text{ w/ } \{\alpha_i, \alpha_j, \alpha_k\} = \{\alpha_1, \alpha_2, \alpha_3\}.$$

We conclude that $S = \emptyset$, i.e. E_λ is smooth.

It is easy to check that Φ is holomorphic and non-constant. To prove it is a biholomorphism,

we only need to show that $\deg \Phi = 1$.

Let us look at $O = [0:1:0]$. It's fibre is

simply $\Phi^{-1}(O) = [0]$. To compute the multiplicity,

consider the chart $z_1 \neq 0$, so that setting

$$x = \frac{z_0}{z_1}, \quad y = \frac{z_2}{z_1},$$

the map Φ is given by $(x, y) = \left(\frac{p(z)}{2p'(z)}, \frac{1}{2p'(z)} \right)$

The curve E_Λ on this chart is $z \left(y - x^3 + \frac{g_2}{4} x y^2 + \frac{g_3}{4} y^3 \right)$

whose derivative in y does not vanish at $(0,0)$. Hence,

it is locally the graph of a function $y(x): \exists U \subseteq \mathbb{C}^2$

neigh of $(0,0)$ and $V \subseteq \mathbb{C}$ neigh. of 0 s.t.

$$V \xrightarrow{\cong} z \left(y - x^3 + \frac{g_2}{4} x y^2 + \frac{g_3}{4} y^3 \right) \cap U$$

$$x \longmapsto (x, y(x))$$

Thus, multiplicity of Φ at $[0]$ is equal to the order

of the zero of $x = \frac{p(z)}{2p'(z)}$ at $z=0$. We know that p

has a simple pole and p' has a triple pole, hence the multiplicity is 1. We conclude that $\deg \Phi = 1$.