

Université de Bordeaux

May 16, 2025

Resurgent large genus asymptotics of intersection numbers

j/w B. Eynard, E. Garcia-Failde, P. Gregori, D. Lewański

arXiv: AG/2309.03143

Alessandro Giacchetto

ETH Zürich

A case study: $m!$

Counting problem:

$$c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:

$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + O(m^{-1}) \right)$$

Con: asymptotically exact

Pro: closed-form

A case study: $m!$

Counting problem:

$$c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:

$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + O(m^{-1}) \right)$$

Con: asymptotically exact

Pro: closed-form

A case study: $m!$

Counting problem:

$$c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:

$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + O(m^{-1}) \right)$$

Con: asymptotically exact

Pro: closed-form

A case study: $m!$

Counting problem: $c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:

$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + O(m^{-1}) \right)$$

Con: asymptotically exact

Pro: closed-form

A case study: $m!$

Counting problem:

$$c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:

$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + O(m^{-1}) \right)$$

Con: asymptotically exact

Pro: closed-form

A case study: $m!$

Counting problem: $c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:

$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + \frac{1}{12} m^{-1} + O(m^{-2}) \right)$$

Con: asymptotically exact

Pro: closed-form

A case study: $m!$

Counting problem: $c_m = \# \left\{ \begin{array}{l} \text{arrangements of } m \text{ distinct objects} \\ \text{into } m \text{ distinct boxes} \end{array} \right\}$

Solution:

$$c_m = m! = \begin{cases} m \cdot c_{m-1} & m > 1 \\ 1 & m = 1 \end{cases}$$

Pro: exact

Con: recursive

Asymptotics:


$$c_m = \sqrt{2\pi m} \left(\frac{m}{e} \right)^m \left(1 + \frac{1}{12} m^{-1} + \frac{1}{288} m^{-2} + O(m^{-3}) \right)$$

Con: asymptotically exact

Pro: closed-form


Visualising Stirling's formula

$$\frac{m!}{\sqrt{2\pi m} \left(\frac{m}{e}\right)^m} - 1 = \mathcal{O}(m^{-1})$$




Visualising Stirling's formula

$$\frac{m!}{\sqrt{2\pi m} \left(\frac{m}{e}\right)^m} - \left(1 + \frac{1}{12}m^{-1}\right) = O(m^{-2})$$



Visualising Stirling's formula

$$\frac{m!}{\sqrt{2\pi m} \left(\frac{m}{e}\right)^m} - \left(1 + \frac{1}{12}m^{-1} + \frac{1}{288}m^{-2}\right) = O(m^{-3})$$



ψ -class intersection numbers

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \quad d_i \geq 0, \quad d_1 + \cdots + d_n = 3g - 3 + n$$

- Volumes of moduli spaces of metric ribbon graphs (maps)

$$V_{g,n}(L_1, \dots, L_n) = \sum_{d_1 + \cdots + d_n = 3g - 3 + n} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \prod_{i=1}^n \frac{L_i^{2d_i}}{2^{d_i} d_i!}$$

- Building block for all tautological intersection numbers:
 - Weil–Petersson volumes
 - Masur–Veech volumes
 - Hurwitz numbers
 - ...
- Compute the perturbative expansion of topological 2d gravity

ψ -class intersection numbers

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \quad d_i \geq 0, \quad d_1 + \cdots + d_n = 3g - 3 + n$$

- Volumes of moduli spaces of **metric ribbon graphs** (maps)

$$V_{g,n}(L_1, \dots, L_n) = \sum_{d_1 + \cdots + d_n = 3g - 3 + n} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \prod_{i=1}^n \frac{L_i^{2d_i}}{2^{d_i} d_i!}$$

- Building block for all **tautological intersection numbers**:
 - Weil–Petersson volumes
 - Masur–Veech volumes
 - Hurwitz numbers
 - ...
- Compute the perturbative expansion of **topological 2d gravity**

ψ -class intersection numbers

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \quad d_i \geq 0, \quad d_1 + \cdots + d_n = 3g - 3 + n$$

- Volumes of moduli spaces of **metric ribbon graphs** (maps)

$$V_{g,n}(L_1, \dots, L_n) = \sum_{d_1 + \cdots + d_n = 3g - 3 + n} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \prod_{i=1}^n \frac{L_i^{2d_i}}{2^{d_i} d_i!}$$

- Building block for all **tautological intersection numbers**:

- Weil–Petersson volumes
 - Masur–Veech volumes
 - Hurwitz numbers
 - ...
- Compute the perturbative expansion of **topological 2d gravity**

ψ -class intersection numbers

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \quad d_i \geq 0, \quad d_1 + \cdots + d_n = 3g - 3 + n$$

- Volumes of moduli spaces of **metric ribbon graphs** (maps)

$$V_{g,n}(L_1, \dots, L_n) = \sum_{d_1 + \cdots + d_n = 3g - 3 + n} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \prod_{i=1}^n \frac{L_i^{2d_i}}{2^{d_i} d_i!}$$

- Building block for all **tautological intersection numbers**:
 - Weil–Petersson volumes
 - Masur–Veech volumes
 - Hurwitz numbers
 - ...
- Compute the perturbative expansion of **topological 2d gravity**

Solution

Normalisation: $\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \prod_{i=1}^n (2d_i + 1)!!$

Witten conjecture/Kontsevich theorem, early '90s:




$$\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = \sum_{m=2}^n (2d_m + 1) \langle\langle \tau_{d_1+d_m-1} \tau_{d_2} \cdots \widehat{\tau_{d_m}} \cdots \tau_{d_n} \rangle\rangle_g$$

$$+ \frac{1}{2} \sum_{a+b=d_1-2} \left(\langle\langle \tau_a \tau_b \tau_{d_2} \cdots \tau_{d_n} \rangle\rangle_{g-1} + \sum_{\substack{g_1+g_2=g \\ I_1 \sqcup I_2 = \{d_2, \dots, d_n\}}} \langle\langle \tau_a \tau_{I_1} \rangle\rangle_{g_1} \langle\langle \tau_b \tau_{I_2} \rangle\rangle_{g_2} \right)$$




with initial data $\langle\langle \tau_0 \tau_0 \tau_0 \rangle\rangle_0 = 1$ and $\langle\langle \tau_1 \rangle\rangle_1 = \frac{1}{8}$.



Solution

Normalisation: $\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \prod_{i=1}^n (2d_i + 1)!!$

Witten conjecture/Kontsevich theorem, early '90s:





$$\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = \sum_{m=2}^n (2d_m + 1) \langle\langle \tau_{d_1+d_m-1} \tau_{d_2} \cdots \widehat{\tau_{d_m}} \cdots \tau_{d_n} \rangle\rangle_g$$

$$+ \frac{1}{2} \sum_{a+b=d_1-2} \left(\langle\langle \tau_a \tau_b \tau_{d_2} \cdots \tau_{d_n} \rangle\rangle_{g-1} + \sum_{\substack{g_1+g_2=g \\ I_1 \sqcup I_2 = \{d_2, \dots, d_n\}}} \langle\langle \tau_a \tau_{I_1} \rangle\rangle_{g_1} \langle\langle \tau_b \tau_{I_2} \rangle\rangle_{g_2} \right)$$



with initial data $\langle\langle \tau_0 \tau_0 \tau_0 \rangle\rangle_0 = 1$ and $\langle\langle \tau_1 \rangle\rangle_1 = \frac{1}{8}$.



Large genus asymptotics

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{\left(\frac{2}{3}\right)^{2g-2+n}} \left(1 + O(g^{-1})\right)$$

- Conjectured by Delecroix–Goujard–Zograf–Zorich, 2019
- Proved by Aggarwal, 2020
(combinatorial/probabilistic analysis of Witten–Kontsevich topological recursion)
- Proved by Guo–Yang, 2021
(combinatorial analysis of the determinantal formula)

Questions

- Universal strategy, adaptable to different problems?
- ‘Geometric’ meaning of the formula?
- Subleading corrections?

Large genus asymptotics

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{\left(\frac{2}{3}\right)^{2g-2+n}} \left(1 + O(g^{-1})\right)$$

- Conjectured by Delecroix–Goujard–Zograf–Zorich, 2019
- Proved by Aggarwal, 2020
(combinatorial/probabilistic analysis of Witten–Kontsevich topological recursion)
- Proved by Guo–Yang, 2021
(combinatorial analysis of the determinantal formula)

Questions

- Universal strategy, adaptable to different problems?
- ‘Geometric’ meaning of the formula?
- Subleading corrections?

Large genus asymptotics

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{\left(\frac{2}{3}\right)^{2g-2+n}} \left(1 + O(g^{-1})\right)$$

- Conjectured by Delecroix–Goujard–Zograf–Zorich, 2019
- Proved by Aggarwal, 2020
(combinatorial/probabilistic analysis of Witten–Kontsevich topological recursion)
- Proved by Guo–Yang, 2021
(combinatorial analysis of the determinantal formula)

Questions

- Universal strategy, adaptable to different problems?
- ‘Geometric’ meaning of the formula?
- Subleading corrections?

Large genus asymptotics: new perspective

Answers

- Universal strategy: resurgent analysis of the determinantal formula
- Geometric meaning: Airy functions

$$y^2 - x = 0 \xrightarrow{\text{quantisation}} \left(\hbar^2 \frac{d^2}{dx^2} - x \right) \psi(x, \hbar) = 0$$

- Subleading corrections: algorithm + properties

Set $(x)_k = x(x-1)\cdots(x-k+1)$.

$$\begin{aligned} \langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} & \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots \right. \\ & \left. + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right) \end{aligned}$$

Large genus asymptotics: new perspective

Answers

- Universal strategy: resurgent analysis of the determinantal formula
- Geometric meaning: Airy functions

$$y^2 - x = 0 \quad \xrightarrow{\text{quantisation}} \quad \left(\hbar^2 \frac{d^2}{dx^2} - x \right) \psi(x, \hbar) = 0$$

- Subleading corrections: algorithm + properties

Set $(x)_k = x(x-1)\cdots(x-k+1)$.

$$\begin{aligned} \langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} & \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots \right. \\ & \left. + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right) \end{aligned}$$

Large genus asymptotics: new perspective

Answers

- Universal strategy: resurgent analysis of the determinantal formula
- Geometric meaning: Airy functions

$$y^2 - x = 0 \xrightarrow{\text{quantisation}} \left(\hbar^2 \frac{d^2}{dx^2} - x \right) \psi(x, \hbar) = 0$$

- Subleading corrections: algorithm + properties

Set $(x)_k = x(x-1)\cdots(x-k+1)$.

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = \underbrace{\frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}}}_{S=1} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

S = 1

Stokes constant

Large genus asymptotics: new perspective

Answers

- Universal strategy: resurgent analysis of the determinantal formula
- Geometric meaning: Airy functions

$$y^2 - x = 0 \xrightarrow{\text{quantisation}} \left(\hbar^2 \frac{d^2}{dx^2} - x \right) \psi(x, \hbar) = 0$$

- Subleading corrections: algorithm + properties

Set $(x)_k = x(x-1)\cdots(x-k+1)$.

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

$$A = 2/3$$

$$\psi \sim \frac{1}{\sqrt{2}x^{1/4}} e^{\pm \frac{A}{\hbar}x^{-3/2}}$$

Large genus asymptotics: new perspective

Answers

- Universal strategy: resurgent analysis of the determinantal formula
- Geometric meaning: Airy functions

$$y^2 - x = 0 \quad \xrightarrow{\text{quantisation}} \quad \left(\hbar^2 \frac{d^2}{dx^2} - x \right) \psi(x, \hbar) = 0$$

- Subleading corrections: algorithm + properties

Set $(x)_k = x(x-1)\cdots(x-k+1)$.

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots \right.$$

$$\left. + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

Computable; polynomial in
 n and multiplicities of d_i

Large genus asymptotics: new perspective

Answers

- Universal strategy: resurgent analysis of the determinantal formula
- Geometric meaning: Airy functions

$$y^2 - x = 0 \xrightarrow{\text{quantisation}} \left(\hbar^2 \frac{d^2}{dx^2} - x \right) \psi(x, \hbar) = 0$$

- Subleading corrections: algorithm + properties

Set $(x)_k = x(x-1)\cdots(x-k+1)$.

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots \right)$$

$$\alpha_1 = -\frac{17-15n+3n^2}{12} - \frac{(3-n)(n-p_0)}{2} - \frac{(n-p_0)_2}{4}$$


$$\text{where } p_0 = \#\{d_i = 0\}$$

$$\frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \Big)$$

Visualising the large genus asymptotics

$$\frac{\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g}{\frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{(2/3)^{2g-2+n}}} = 1 + O(g^{-1})$$


For $n = 2$:



Visualising the large genus asymptotics

$$\frac{2g-3+n}{2/3} \left(\frac{\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g}{\frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{(2/3)^{2g-2+n}}} - 1 \right) = \alpha_1(n, p_0) + O(g^{-1})$$

For $n = 2$:



Borel meets Darboux

Borel's idea:

- Divergent power series:

$$\tilde{\varphi}(\hbar) = \sum_{m \geq 0} a_m \hbar^m$$

with $|a_m| = O(R^{-m} m!)$.

- The **Borel transform**

$$\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$$

is now abs. convergent

Darboux's idea:

- Abs. convergent power series:

$$\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$$

- Get a holomorphic function around the origin, take analytic continuation

- The large m **asymptotics** of a_m is totally controlled by the behaviour of $\hat{\varphi}$ at its **singularities**

Borel meets Darboux

Borel's idea:

- Divergent power series:

$$\tilde{\varphi}(\hbar) = \sum_{m \geq 0} a_m \hbar^m$$

with $|a_m| = O(R^{-m} m!)$.

- The **Borel transform**

$$\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$$

is now abs. convergent

Darboux's idea:

- Abs. convergent power series:

$$\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$$

- Get a holomorphic function around the origin, take analytic continuation

- The large m **asymptotics** of a_m is totally controlled by the behaviour of $\hat{\varphi}$ at its **singularities**

Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$

Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = (\text{holomorphic @ } A) \log(s - A) + \text{holomorphic @ } A$$

$$a_m = \frac{m!}{2\pi i} \oint_C \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$




Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$

Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = (\text{holomorphic @ } A) \log(s - A) + \text{holomorphic @ } A$$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$




Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$

Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$




Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$

Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof


Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$

Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

Stokes constant
 $s \in \mathbb{C}$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic } @ A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$a_m = \frac{m!}{2\pi i} \oint_{\mathcal{C}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$a_m = \frac{m!}{2\pi i} \int_{\mathcal{H}} \frac{\hat{\varphi}(s)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic } @ A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$a_m = -\frac{m!}{2\pi i} \int_{\mathcal{H}} \frac{s}{2\pi} \frac{\hat{\psi}(s-A)}{s^{m+1}} \log(s-A) ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$a_m = \frac{s}{2\pi} m! \int_A^\infty \frac{\hat{\psi}(s-A)}{s^{m+1}} ds$$



Darboux's result: sketch of the proof

Take an abs. convergent power series: $\hat{\varphi}(s) = \sum_{m \geq 0} \frac{a_m}{m!} s^m$


Suppose its analytic continuation has a single log singularity at $s = A$:

$$\hat{\varphi}(s) = -\frac{s}{2\pi} \hat{\psi}(s-A) \log(s-A) + \text{holomorphic @ } A$$

Stokes constant
 $s \in \mathbb{C}$

minor
holom @ origin
 $\hat{\psi}(s) = \sum_{k \geq 0} \frac{b_k}{k!} s^k$

$$\begin{aligned} a_m &= \frac{s}{2\pi} \frac{\Gamma(m)}{A^m} \left(b_0 + \frac{A}{m-1} b_1 + \dots \right. \\ &\quad \left. + \frac{A^k}{(m-1)_k} b_k + O(m^{-k-1}) \right) \end{aligned}$$



Borel meets Darboux: the algorithm

- Given: $\tilde{\varphi}(\hbar) = \sum_m a_m \hbar^m$ divergent
- Borel transform: $\hat{\varphi}(s) = \sum_m \frac{a_m}{m!} s^m$ abs. convergent
- Suppose you can compute:
 - Log singularities: $A \in \text{Sing}(\hat{\varphi})$
 - Stokes constants: $(S_A)_{A \in \text{Sing}(\hat{\varphi})}$
 - Minors: $(\hat{\Psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$
- Large m asymptotics:

$$a_m = \frac{\Gamma(m)}{2\pi} \sum_{A \in \text{Sing}(\hat{\varphi})} \frac{S_A}{A^m} \left(b_{A,0} + \frac{A}{m-1} b_{A,1} + \frac{A^2}{(m-1)(m-2)} b_{A,2} + \dots \right)$$

Borel meets Darboux: the algorithm

- Given: $\tilde{\varphi}(\hbar) = \sum_m a_m \hbar^m$ divergent
- Borel transform: $\hat{\varphi}(s) = \sum_m \frac{a_m}{m!} s^m$ abs. convergent
- Suppose you can compute:
 - Log singularities: $A \in \text{Sing}(\hat{\varphi})$
 - Stokes constants: $(S_A)_{A \in \text{Sing}(\hat{\varphi})}$
 - Minors: $(\hat{\Psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$
- Large m asymptotics:

$$a_m = \frac{\Gamma(m)}{2\pi} \sum_{A \in \text{Sing}(\hat{\varphi})} \frac{S_A}{A^m} \left(b_{A,0} + \frac{A}{m-1} b_{A,1} + \frac{A^2}{(m-1)(m-2)} b_{A,2} + \dots \right)$$

Borel meets Darboux: the algorithm

- Given: $\tilde{\varphi}(\hbar) = \sum_m a_m \hbar^m$ divergent
- Borel transform: $\hat{\varphi}(s) = \sum_m \frac{a_m}{m!} s^m$ abs. convergent
- Suppose you can compute:
 - ① Log **singularities**: $A \in \text{Sing}(\hat{\varphi})$
 - ② **Stokes constants**: $(S_A)_{A \in \text{Sing}(\hat{\varphi})}$
 - ③ **Minors**: $(\hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$
- Large m **asymptotics**:

$$a_m = \frac{\Gamma(m)}{2\pi} \sum_{A \in \text{Sing}(\hat{\varphi})} \frac{S_A}{A^m} \left(b_{A,0} + \frac{A}{m-1} b_{A,1} + \frac{A^2}{(m-1)(m-2)} b_{A,2} + \dots \right)$$

Borel meets Darboux: the algorithm

- Given: $\tilde{\varphi}(\hbar) = \sum_m a_m \hbar^m$ divergent
- Borel transform: $\hat{\varphi}(s) = \sum_m \frac{a_m}{m!} s^m$ abs. convergent
- Suppose you can compute:
 - ① Log **singularities**: $A \in \text{Sing}(\hat{\varphi})$
 - ② **Stokes constants**: $(S_A)_{A \in \text{Sing}(\hat{\varphi})}$
 - ③ **Minors**: $(\hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$
- Large m **asymptotics**:

$$a_m = \frac{\Gamma(m)}{2\pi} \sum_{A \in \text{Sing}(\hat{\varphi})} \frac{S_A}{A^m} \left(b_{A,0} + \frac{A}{m-1} b_{A,1} + \frac{A^2}{(m-1)(m-2)} b_{A,2} + \dots \right)$$

Properties of the resurgence method

- **Algorithmic.**

$$\tilde{\varphi} = \sum_m a_m \hbar^m \quad \rightarrow \quad (S_A, \hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})} \quad \rightarrow \quad \text{asymptotic of } a_m$$

- **Exponential integrals.** The singularity structure of exponential integrals is well-understood:

$$\tilde{\varphi} = \text{Asym} \left(\int e^{-\frac{1}{\hbar} S(t)} dt \right) \quad \rightarrow \quad (S_A, \hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$$

- **Sums and products.** The singularity structure of sums and products of divergent series is well-understood:

$$\begin{aligned} \lambda_1 \tilde{\varphi}_1 + \lambda_2 \tilde{\varphi}_2 &\quad \rightarrow \quad (S_A^+, \hat{\psi}_A^+)_{A \in \text{Sing}(\hat{\varphi}_1) \cup \text{Sing}(\hat{\varphi}_2)} \\ \tilde{\varphi}_1 \cdot \tilde{\varphi}_2 &\quad \rightarrow \quad (S_A^\times, \hat{\psi}_A^\times)_{A \in \text{Sing}(\hat{\varphi}_1) \cup \text{Sing}(\hat{\varphi}_2)} \end{aligned}$$

Properties of the resurgence method

- **Algorithmic.**

$$\tilde{\varphi} = \sum_m a_m \hbar^m \quad \rightarrow \quad (S_A, \hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})} \quad \rightarrow \quad \text{asymptotic of } a_m$$

- **Exponential integrals.** The singularity structure of exponential integrals is well-understood:

$$\tilde{\varphi} = \text{Asym} \left(\int e^{-\frac{1}{\hbar} S(t)} dt \right) \quad \rightarrow \quad (S_A, \hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$$

- **Sums and products.** The singularity structure of sums and products of divergent series is well-understood:

$$\begin{aligned} \lambda_1 \tilde{\varphi}_1 + \lambda_2 \tilde{\varphi}_2 &\quad \rightarrow \quad (S_A^+, \hat{\psi}_A^+)_{A \in \text{Sing}(\hat{\varphi}_1) \cup \text{Sing}(\hat{\varphi}_2)} \\ \tilde{\varphi}_1 \cdot \tilde{\varphi}_2 &\quad \rightarrow \quad (S_A^\times, \hat{\psi}_A^\times)_{A \in \text{Sing}(\hat{\varphi}_1) \cup \text{Sing}(\hat{\varphi}_2)} \end{aligned}$$

Properties of the resurgence method

- **Algorithmic.**

$$\tilde{\varphi} = \sum_m a_m \hbar^m \quad \longrightarrow \quad (S_A, \hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})} \quad \longrightarrow \quad \text{asymptotic of } a_m$$

- **Exponential integrals.** The singularity structure of exponential integrals is well-understood:

$$\tilde{\varphi} = \text{Asym} \left(\int e^{-\frac{1}{\hbar} S(t)} dt \right) \quad \longrightarrow \quad (S_A, \hat{\psi}_A)_{A \in \text{Sing}(\hat{\varphi})}$$

- **Sums and products.** The singularity structure of sums and products of divergent series is well-understood:

$$\begin{aligned} \lambda_1 \tilde{\varphi}_1 + \lambda_2 \tilde{\varphi}_2 &\quad \longrightarrow \quad (S_A^+, \hat{\psi}_A^+)_{A \in \text{Sing}(\hat{\varphi}_1) \cup \text{Sing}(\hat{\varphi}_2)} \\ \tilde{\varphi}_1 \cdot \tilde{\varphi}_2 &\quad \longrightarrow \quad (S_A^\times, \hat{\psi}_A^\times)_{A \in \text{Sing}(\hat{\varphi}_1) \cup \text{Sing}(\hat{\varphi}_2)} \end{aligned}$$

Example: the Airy kernel

$$\begin{aligned}
 K(z, w; \hbar) &= \frac{\text{Ai}(z^2; \hbar)\text{Bi}'(w^2; \hbar) - \text{Ai}'(z^2; \hbar)\text{Bi}(w^2; \hbar)}{z^2 - w^2} = \sum_{m \geq 0} a_m \hbar^m \\
 &= \frac{1}{2\sqrt{zw(z-w)}} - \frac{1}{(zw)^{3/2}} \left(\frac{5}{96z^2} - \frac{7}{96zw} + \frac{5}{96w^2} \right) \hbar \\
 &\quad + \frac{1}{(zw)^{3/2}} \left(\frac{385}{9216z^5} - \frac{455}{9216z^4w} + \frac{385}{9216z^3w^2} - \frac{385}{9216z^2w^3} + \frac{455}{9216zw^4} - \frac{385}{9216w^5} \right) \hbar^2 + \dots
 \end{aligned}$$

$\widehat{\text{Ai}}(z^2; \hbar)$ and $\widehat{\text{Ai}'}(z^2; \hbar)$ have

- a **log singularity** at $\frac{4}{3}z^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Bi}(z^2; \hbar)$ and $\text{Bi}'(z^2; \hbar)$

$\widehat{\text{Bi}}(w^2; \hbar)$ and $\widehat{\text{Bi}'}(w^2; \hbar)$ have

- a **log singularity** at $-\frac{4}{3}w^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Ai}(w^2; \hbar)$ and $\text{Ai}'(w^2; \hbar)$

$$\Rightarrow a_m = \frac{\Gamma(m)}{2\pi} \left(\frac{1}{(\frac{4}{3}z^3)^m} \frac{w-z}{2\sqrt{zw}(z^2-w^2)} + \frac{1}{(-\frac{4}{3}w^3)^m} \frac{z-w}{2\sqrt{zw}(z^2-w^2)} + \dots \right)$$

Example: the Airy kernel

$$\begin{aligned}
 K(z, w; \hbar) &= \frac{\text{Ai}(z^2; \hbar)\text{Bi}'(w^2; \hbar) - \text{Ai}'(z^2; \hbar)\text{Bi}(w^2; \hbar)}{z^2 - w^2} = \sum_{m \geq 0} a_m \hbar^m \\
 &= \frac{1}{2\sqrt{zw}(z-w)} - \frac{1}{(zw)^{3/2}} \left(\frac{5}{96z^2} - \frac{7}{96zw} + \frac{5}{96w^2} \right) \hbar \\
 &\quad + \frac{1}{(zw)^{3/2}} \left(\frac{385}{9216z^5} - \frac{455}{9216z^4w} + \frac{385}{9216z^3w^2} - \frac{385}{9216z^2w^3} + \frac{455}{9216zw^4} - \frac{385}{9216w^5} \right) \hbar^2 + \dots
 \end{aligned}$$

$\widehat{\text{Ai}}(z^2; \hbar)$ and $\widehat{\text{Ai}'}(z^2; \hbar)$ have

- a **log singularity** at $\frac{4}{3}z^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Bi}(z^2; \hbar)$ and $\text{Bi}'(z^2; \hbar)$

$\widehat{\text{Bi}}(w^2; \hbar)$ and $\widehat{\text{Bi}'}(w^2; \hbar)$ have

- a **log singularity** at $-\frac{4}{3}w^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Ai}(w^2; \hbar)$ and $\text{Ai}'(w^2; \hbar)$

$$\Rightarrow a_m = \frac{\Gamma(m)}{2\pi} \left(\frac{1}{(\frac{4}{3}z^3)^m} \frac{w-z}{2\sqrt{zw}(z^2-w^2)} + \frac{1}{(-\frac{4}{3}w^3)^m} \frac{z-w}{2\sqrt{zw}(z^2-w^2)} + \dots \right)$$

Example: the Airy kernel

$$\begin{aligned}
 K(z, w; \hbar) &= \frac{\text{Ai}(z^2; \hbar)\text{Bi}'(w^2; \hbar) - \text{Ai}'(z^2; \hbar)\text{Bi}(w^2; \hbar)}{z^2 - w^2} = \sum_{m \geq 0} a_m \hbar^m \\
 &= \frac{1}{2\sqrt{zw(z-w)}} - \frac{1}{(zw)^{3/2}} \left(\frac{5}{96z^2} - \frac{7}{96zw} + \frac{5}{96w^2} \right) \hbar \\
 &\quad + \frac{1}{(zw)^{3/2}} \left(\frac{385}{9216z^5} - \frac{455}{9216z^4w} + \frac{385}{9216z^3w^2} - \frac{385}{9216z^2w^3} + \frac{455}{9216zw^4} - \frac{385}{9216w^5} \right) \hbar^2 + \dots
 \end{aligned}$$

$\widehat{\text{Ai}}(z^2; \hbar)$ and $\widehat{\text{Ai}}'(z^2; \hbar)$ have

- a **log singularity** at $\frac{4}{3}z^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Bi}(z^2; \hbar)$ and $\text{Bi}'(z^2; \hbar)$

$\widehat{\text{Bi}}(w^2; \hbar)$ and $\widehat{\text{Bi}}'(w^2; \hbar)$ have

- a **log singularity** at $-\frac{4}{3}w^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Ai}(w^2; \hbar)$ and $\text{Ai}'(w^2; \hbar)$

$$\Rightarrow a_m = \frac{\Gamma(m)}{2\pi} \left(\frac{1}{(\frac{4}{3}z^3)^m} \frac{w-z}{2\sqrt{zw(z^2-w^2)}} + \frac{1}{(-\frac{4}{3}w^3)^m} \frac{z-w}{2\sqrt{zw(z^2-w^2)}} + \dots \right)$$

Example: the Airy kernel

$$\begin{aligned}
 K(z, w; \hbar) &= \frac{\text{Ai}(z^2; \hbar)\text{Bi}'(w^2; \hbar) - \text{Ai}'(z^2; \hbar)\text{Bi}(w^2; \hbar)}{z^2 - w^2} = \sum_{m \geq 0} a_m \hbar^m \\
 &= \frac{1}{2\sqrt{zw}(z-w)} - \frac{1}{(zw)^{3/2}} \left(\frac{5}{96z^2} - \frac{7}{96zw} + \frac{5}{96w^2} \right) \hbar \\
 &\quad + \frac{1}{(zw)^{3/2}} \left(\frac{385}{9216z^5} - \frac{455}{9216z^4w} + \frac{385}{9216z^3w^2} - \frac{385}{9216z^2w^3} + \frac{455}{9216zw^4} - \frac{385}{9216w^5} \right) \hbar^2 + \dots
 \end{aligned}$$

$\widehat{\text{Ai}}(z^2; \hbar)$ and $\widehat{\text{Ai}'}(z^2; \hbar)$ have

- a **log singularity** at $\frac{4}{3}z^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Bi}(z^2; \hbar)$ and $\text{Bi}'(z^2; \hbar)$

$\widehat{\text{Bi}}(w^2; \hbar)$ and $\widehat{\text{Bi}'}(w^2; \hbar)$ have


- a **log singularity** at $-\frac{4}{3}w^3$
- **Stokes** constant: $S = 1$
- **minors**: $\text{Ai}(w^2; \hbar)$ and $\text{Ai}'(w^2; \hbar)$

$$\implies a_m = \frac{(-1)^m}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m} \left(\frac{1}{(zw)^{3/2}} h_{3m-1}\left(\frac{1}{z}, -\frac{1}{w}\right) + \dots \right)$$

Example visualised

Write $a_m = \frac{(-1)^m}{(zw)^{3/2}} \sum_{k+\ell=3m-1} a_{k,\ell} \frac{1}{z^k (-w)^\ell}$.


$$a_{k,\ell} = \frac{1}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m} \left(1 + O(m^{-1})\right) \implies \frac{a_{k,\ell}}{\frac{1}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m}} = 1 + O(m^{-1})$$



Example visualised

Write $a_m = \frac{(-1)^m}{(zw)^{3/2}} \sum_{k+\ell=3m-1} a_{k,\ell} \frac{1}{z^k (-w)^\ell}$.


$$a_{k,\ell} = \frac{1}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m} \left(1 + O(m^{-1})\right) \implies \frac{a_{k,\ell}}{\frac{1}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m}} = 1 + O(m^{-1})$$



Example visualised

Write $a_m = \frac{(-1)^m}{(zw)^{3/2}} \sum_{k+\ell=3m-1} a_{k,\ell} \frac{1}{z^k (-w)^\ell}$.

$$a_{k,\ell} = \frac{1}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m} \left(1 + O(m^{-1})\right) \implies \frac{a_{k,\ell}}{\frac{1}{4\pi} \frac{\Gamma(m)}{\left(\frac{4}{3}\right)^m}} = 1 + O(m^{-1})$$



Strategy towards large genus asymptotics

Goal

Compute the large genus asymptotics of $\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g$

- ① Take the n -pnt fnct

$$W_n(z_1, \dots, z_n; \hbar) = \sum_{g \geq 0} \hbar^{2g-2+n} W_{g,n}(z_1, \dots, z_n)$$

- ② W_n is a divergent series in \hbar . Take its Borel transform and study its singularity structure
- ③ Get the large genus asymptotics (with subleading contributions!)

Strategy towards large genus asymptotics

Goal

Compute the large genus asymptotics of $\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g$

- ➊ Take the n -pnt fnct

$$W_n(z_1, \dots, z_n; \hbar) = \sum_{g \geq 0} \hbar^{2g-2+n} W_{g,n}(z_1, \dots, z_n)$$

- ➋ W_n is a divergent series in \hbar . Take its Borel transform and study its singularity structure
- ➌ Get the large genus asymptotics (with subleading contributions!)

Strategy towards large genus asymptotics

Goal

Compute the large genus asymptotics of $\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g$

- ① Take the n -pnt fnct

$$\begin{aligned} W_n(z_1, \dots, z_n; \hbar) &= \sum_{g \geq 0} \hbar^{2g-2+n} W_{g,n}(z_1, \dots, z_n) \\ &= (-2)^{-(2g-2+n)} \sum_{d_1, \dots, d_n} \frac{\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g}{2z_1^{2d_1+3} \cdots 2z_n^{2d_n+3}} \end{aligned}$$

- ② W_n is a divergent series in \hbar . Take its Borel transform and study its singularity structure
- ③ Get the large genus asymptotics (with subleading contributions!)

Strategy towards large genus asymptotics

Goal

Compute the large genus asymptotics of $\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g$

- ① Take the n -pt fnct

$$\begin{aligned} W_n(z_1, \dots, z_n; \hbar) &= \sum_{g \geq 0} \hbar^{2g-2+n} W_{g,n}(z_1, \dots, z_n) \\ &= (-2)^{-(2g-2+n)} \sum_{d_1, \dots, d_n} \frac{\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g}{2z_1^{2d_1+3} \cdots 2z_n^{2d_n+3}} \end{aligned}$$

n fixed
 $\hbar \rightarrow$ genus
 $z_i \rightarrow d_i$

- ② W_n is a divergent series in \hbar . Take its Borel transform and study its singularity structure
- ③ Get the large genus asymptotics (with subleading contributions!)

Strategy towards large genus asymptotics

Goal

Compute the large genus asymptotics of $\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g$

- ① Take the n -pt fnct

n fixed
 $\hbar \rightarrow$ genus
 $z_i \rightarrow d_i$

$$\begin{aligned} W_n(z_1, \dots, z_n; \hbar) &= \sum_{g \geq 0} \hbar^{2g-2+n} W_{g,n}(z_1, \dots, z_n) \\ &= (-2)^{-(2g-2+n)} \sum_{d_1, \dots, d_n} \frac{\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g}{2z_1^{2d_1+3} \cdots 2z_n^{2d_n+3}} \end{aligned}$$

- ② W_n is a divergent series in \hbar . Take its Borel transform and study its singularity structure
- ③ Get the large genus asymptotics (with subleading contributions!)

Strategy towards large genus asymptotics

Goal

Compute the large genus asymptotics of $\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g$

- 1 Take the n -pt fnct

$$\begin{aligned} W_n(z_1, \dots, z_n; \hbar) &= \sum_{g \geq 0} \hbar^{2g-2+n} W_{g,n}(z_1, \dots, z_n) \\ &= (-2)^{-(2g-2+n)} \sum_{d_1, \dots, d_n} \frac{\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g}{2z_1^{2d_1+3} \cdots 2z_n^{2d_n+3}} \end{aligned}$$

n fixed
 $\hbar \rightarrow$ genus
 $z_i \rightarrow d_i$

- 2 W_n is a divergent series in \hbar . Take its Borel transform and study its singularity structure
- 3 Get the large genus asymptotics (with subleading contributions!)

Determinantal formula

Define the disconnected n -pnt fnct and recall the Airy kernel

$$W_n^\bullet(z_1, \dots, z_n; \hbar) = \sum_{P \in \text{Part}(n)} W_{\ell(P)}(z_P; \hbar),$$

$$K(z, w; \hbar) = \frac{\text{Ai}(z^2; \hbar)\text{Bi}'(w^2; \hbar) - \text{Ai}'(z^2; \hbar)\text{Bi}(w^2; \hbar)}{z^2 - w^2}.$$

Determinantal formula (Bergère–Eynard, Bertola–Dubrovin–Yang):

$$W_n^\bullet(z_1, \dots, z_n; \hbar) = \det_{1 \leqslant i, j \leqslant n} K(z_i, z_j; \hbar)$$

Determinantal formula

Define the disconnected n -pnt fnct and recall the Airy kernel

$$W_n^\bullet(z_1, \dots, z_n; \hbar) = \sum_{P \in \text{Part}(n)} W_{\ell(P)}(z_P; \hbar),$$

$$K(z, w; \hbar) = \frac{\text{Ai}(z^2; \hbar)\text{Bi}'(w^2; \hbar) - \text{Ai}'(z^2; \hbar)\text{Bi}(w^2; \hbar)}{z^2 - w^2}.$$

Determinantal formula (Bergère–Eynard, Bertola–Dubrovin–Yang):

$$W_n^\bullet(z_1, \dots, z_n; \hbar) = \det_{1 \leqslant i, j \leqslant n} K(z_i, z_j; \hbar)$$

Singularity structure of W_n

Singularity strct
of A_i, B_i



Singularity strct
of W_n

- $2n \log$ singularities of \widehat{W}_n , located at

$$+ \frac{4}{3}z_i^3 \quad \text{and} \quad - \frac{4}{3}z_i^3, \quad i = 1, \dots, n$$

- Stokes constants: $S = 1$

- Minors:

(A) at $+ \frac{4}{3}z_i^3$: replace each (A_i, A'_i) with (B_i, B'_i)

(B) at $- \frac{4}{3}z_i^3$: replace each (B_i, B'_i) with (A_i, A'_i)

Singularity structure of W_n

Singularity strct
of A_i, B_i



Singularity strct
of W_n

- $2n \log$ singularities of \widehat{W}_n , located at

$$+ \frac{4}{3}z_i^3 \quad \text{and} \quad - \frac{4}{3}z_i^3, \quad i = 1, \dots, n$$

- Stokes constants: $S = 1$
- Minors:

(A) at $+ \frac{4}{3}z_i^3$: replace each (A_i, A'_i) with (B_i, B'_i)

(B) at $- \frac{4}{3}z_i^3$: replace each (B_i, B'_i) with (A_i, A'_i)

Singularity structure of W_n

Singularity strct
of A_i, B_i



Singularity strct
of W_n

- $2n \log$ singularities of \widehat{W}_n , located at

$$+ \frac{4}{3}z_i^3 \quad \text{and} \quad - \frac{4}{3}z_i^3, \quad i = 1, \dots, n$$

- Stokes constants: $S = 1$

- Minors:

(A) at $+ \frac{4}{3}z_i^3$: replace each (A_i, A'_i) with (B_i, B'_i)

(B) at $- \frac{4}{3}z_i^3$: replace each (B_i, B'_i) with (A_i, A'_i)

Singularity structure of W_n

Singularity strct
of A_i, B_i



Singularity strct
of W_n

- $2n \log$ singularities of \widehat{W}_n , located at

$$+ \frac{4}{3}z_i^3 \quad \text{and} \quad - \frac{4}{3}z_i^3, \quad i = 1, \dots, n$$

- Stokes constants: $S = 1$

- Minors:

A at $+ \frac{4}{3}z_i^3$: replace each (A_i, A'_i) with (B_i, B'_i)

B at $- \frac{4}{3}z_i^3$: replace each (B_i, B'_i) with (A_i, A'_i)

Summary

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\!\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\!\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

where:

- $S = 1$
Stokes constants of the Airy ODE
- $A = 2/3$
leading exp behaviour of Ai
- α_k polynomials in n and multiplicities of d_i (conj by Guo-Yang)
are computable from the asymptotic expansion coeffs of Ai

Summary

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

where:

- $S = 1$
Stokes constants of the Airy ODE
- $A = 2/3$
leading exp behaviour of A_i
- α_k polynomials in n and multiplicities of d_i (conj by Guo-Yang)
are computable from the asymptotic expansion coeffs of A_i

Summary

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

where:

- $S = 1$
Stokes constants of the Airy ODE
- $A = 2/3$
leading exp behaviour of Ai
- α_k polynomials in n and multiplicities of d_i (conj by Guo-Yang)
are computable from the asymptotic expansion coeffs of Ai

Summary

Uniformly in d_1, \dots, d_n as $g \rightarrow \infty$:

$$\langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g = S \frac{2^n}{4\pi} \frac{\Gamma(2g-2+n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g-3+n} \alpha_1 + \cdots + \frac{A^k}{(2g-3+n)_k} \alpha_k + O(g^{-k-1}) \right)$$

where:

- $S = 1$
Stokes constants of the Airy ODE
- $A = 2/3$
leading exp behaviour of Ai
- α_k polynomials in n and multiplicities of d_i (conj by Guo-Yang)
are computable from the asymptotic expansion coeffs of Ai

Bessel

Norbury's int. nmbrs (BGW τ -fnct (Chidambaram–Garcia-Failde–AG)):

$$\begin{aligned} \langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g^\Theta &= \int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \prod_{i=1}^n \psi_i^{d_i} (2d_i + 1)!! \\ &= S \frac{2^n}{4\pi} \frac{\Gamma(2g - 2 + n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g - 3 + n} \alpha_1 + \cdots \right. \\ &\quad \left. + \frac{A^k}{(2g - 3 + n)^k} \alpha_k + O(g^{-k-1}) \right) \end{aligned}$$

where:

- $S = 2$
Stokes constants of the Bessel ODE
- $A = 2$
leading exp behaviour of K_0
- α_k polynomials in n and multiplicities of d_i
are computable from the asymptotic expansion coeffs of K_0

Bessel

Norbury's int. nmbrs (BGW τ -fnct (Chidambaram–Garcia-Failde–AG)):

$$\begin{aligned} \langle\langle \tau_{d_1} \cdots \tau_{d_n} \rangle\rangle_g^\Theta &= \int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \prod_{i=1}^n \psi_i^{d_i} (2d_i + 1)!! \\ &= S \frac{2^n}{4\pi} \frac{\Gamma(2g - 2 + n)}{A^{2g-2+n}} \left(1 + \frac{A}{2g - 3 + n} \alpha_1 + \cdots \right. \\ &\quad \left. + \frac{A^k}{(2g - 3 + n)^k} \alpha_k + O(g^{-k-1}) \right) \end{aligned}$$

where:

- $S = 2$

Stokes constants of the Bessel ODE

- $A = 2$

leading exp behaviour of K_0

- α_k polynomials in n and multiplicities of d_i

are computable from the asymptotic expansion coeffs of K_0

r-Airy

Witten *r*-spin int. nmbrs (*r*-KdV τ -fnct (Faber–Shadrin–Zvonkine)):

$$\begin{aligned}
 \langle\!\langle \tau_{a_1, d_1} \cdots \tau_{a_n, d_n} \rangle\!\rangle_g^{r\text{-spin}} &= \int_{\overline{\mathcal{M}}_{g,n}} c_w(a_1, \dots, a_n) \prod_{i=1}^n \psi_i^{d_i} (rd_i + a_i)!_{(r)} \\
 &= \frac{2^n}{2\pi} \frac{\Gamma(2g-2+n)}{r^{g-1-|d|}} \left[\frac{S_{r,1}}{|A_{r,1}|^{2g-2+n}} \left(\alpha_0^{(r,1)} + \frac{|A_{r,1}|}{2g-3+n} \alpha_1^{(r,1)} + \dots \right) \right. \\
 &\quad + \dots \\
 &\quad + \frac{S_{r,\lfloor \frac{r-1}{2} \rfloor}}{|A_{r,\lfloor \frac{r-1}{2} \rfloor}|^{2g-2+n}} \left(\alpha_0^{(r,\lfloor \frac{r-1}{2} \rfloor)} + \frac{|A_{r,\lfloor \frac{r-1}{2} \rfloor}|^K}{2g-3+n} \alpha_1^{(r,\lfloor \frac{r-1}{2} \rfloor)} + \dots \right) \\
 &\quad \left. + \frac{\delta_r^{\text{even}}}{2} \frac{S_{r,\frac{r}{2}}}{|A_{r,\frac{r}{2}}|^{2g-2+n}} \left(\alpha_0^{(r,\frac{r}{2})} + \frac{|A_{r,\frac{r}{2}}|^K}{2g-3+n} \alpha_1^{(r,\frac{r}{2})} + \dots \right) \right]
 \end{aligned}$$

where $S_{r,i}$, $A_{r,i}$, $\alpha_k^{(r,i)}$ are obtained the *r*-Airy ODE.

Thank you for the attention!